Serial No.: 10/613,228 - 2 -

Conf. No.: 4680

In the Claims

Please replace all prior versions, and listings, of claims in the application with the following list of claims:

1. (Original) A composition comprising

an immunostimulatory nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1.

- 2. (Original) The composition of claim 1, wherein the immunostimulatory nucleic acid molecule consists of the nucleotide sequence of SEQ ID NO:1.
 - 3. (Original) The composition of claim 1, further comprising an antigen.
- 4. (Original) The composition of claim 3, wherein the antigen is selected from the group consisting of a microbial antigen, a cancer antigen, and an allergen.
- 5. (Original) The composition of claim 4, wherein the microbial antigen is selected from the group consisting of a bacterial antigen, a viral antigen, a fungal antigen and a parasitic antigen.
- 6. (Original) The composition of claim 3, wherein the antigen is encoded by a nucleic acid vector.
- 7. (Original) The composition of claim 3, wherein the nucleic acid vector is separate from the immunostimulatory nucleic acid.
 - 8. (Original) The composition of claim 3, wherein the antigen is a peptide antigen.
 - 9. (Original) The composition of claim 1, further comprising an adjuvant.
 - 10. (Original) The composition of claim 9, wherein the adjuvant is a mucosal adjuvant.

Serial No.: 10/613,228 - 3 -

Conf. No.: 4680

11. (Original) The composition of claim 1, further comprising a cytokine.

12. (Currently Amended) The composition of claim 1, further comprising a therapeutic agent selected from the group consisting of an anti-microbial agent, an anti-cancer agent, <u>and</u> an allergy/asthma medicament.

- 13. (Original) The composition of claim 12, wherein the anti-microbial agent is selected from the group consisting of an anti-bacterial agent, an anti-viral agent, an anti-fungal agent, and an anti-parasite agent.
- 14. (Original) The composition of claim 12, wherein the anti-cancer agent is selected from the group consisting of a chemotherapeutic agent, a cancer vaccine, and an immunotherapeutic agent.
- 15. (Original) The composition of claim 12, wherein the allergy/asthma medicament is selected from the group consisting of PDE-4 inhibitor, bronchodilator/beta-2 agonist, K+ channel opener, VLA-4 antagonist, neurokin antagonist, TXA2 synthesis inhibitor, xanthanine, arachidonic acid antagonist, 5 lipoxygenase inhibitor, thromboxin A2 receptor antagonist, thromboxane A2 antagonist, inhibitor of 5-lipox activation protein, and protease inhibitor.
- 17. (Original) The composition of claim 1, wherein the immunostimulatory nucleic acid has a nucleotide backbone which includes at least one backbone modification.
- 18. (Original) The composition of claim 17, wherein the backbone modification is a phosphorothicate modification.
- 19. (Original) The composition of claim 17, wherein the nucleotide backbone is chimeric.

Serial No.: 10/613,228 - 4 -

Conf. No.: 4680

20. (Original) The composition of claim 17, wherein the nucleotide backbone is entirely modified.

21. (Original) The composition of claim 1, further comprising a pharmaceutically acceptable carrier.

22. (Cancelled)

23. (Original) The composition of claim 1, wherein the immunostimulatory nucleic acid includes at least four CpG motifs.

24.-27. (Cancelled)

- 28. (Original) The composition of claim 1, wherein the immunostimulatory nucleic acid is formulated as a nutritional supplement.
- 29. (Original) The composition of claim 28, wherein the nutritional supplement is formulated as a capsule, a pill, or a sublingual tablet.
- 30. (Original) The composition of claim 1, wherein the immunostimulatory nucleic acid is formulated for local administration.
- 31. (Original) The c omposition of claim 1, wherein the immunostimulatory nucleic acid is formulated for parenteral administration.
- 32. (Original) The composition of claim 1, wherein the immunostimulatory nucleic acid is formulated in a sustained release device.
- 33. (Original) The composition of claim 1, wherein the immunostimulatory nucleic acid is formulated for delivery to a mucosal surface.

Serial No.: 10/613,228 - 5 -

Conf. No.: 4680

34.-43. (Cancelled)

44. (Original) The composition of claim 32, wherein the sustained release device is a microparticle.

45. (Cancelled)

46. (Currently Amended) A method for stimulating an immune response in a subject in need thereof comprising

administering to a subject an the immunostimulatory nucleic acid molecule emprising the nucleotide sequence of SEQ ID NO:1 of claim 1, in an amount effective to stimulate an immune response.

- 47. (Original) The method of claim 46, wherein the subject has or is at risk of developing an infection.
- 48. (Original) The method of claim 47, wherein the infection is selected from the group consisting of a bacterial infection, a viral infection, a fungal infection, and a parasite infection.
- 49. (Original) The method of claim 48, wherein the viral infection is selected from the group consisting of Human immunodeficiency viruses (HIV-1 and HIV-2), Human T lymphotropic virus type I (HTLV-I), Human T lymphotrophic virus type II (HTLV-II), Herpes simplex virus type I (HSV-1) Herpes simplex virus type 2 (HSV-2), Human papilloma virus (multiple types), Hepatitis A virus, Hepatitis B virus, Hepatitis C and D viruses, Epstein-Barr virus (EBV), Cytomegalovirus and Molluscum contagiosum virus.
- 50. (Original) The method of claim 49, wherein the viral infection is a herpes simplex virus infection.
- 51. (Original) The method of claim 46, wherein the subject has or is at risk of developing allergy.

Serial No.: 10/613,228 - 6 -

Conf. No.: 4680

52. (Original) The method of claim 46, wherein the subject has or is at risk of developing asthma.

- 53. (Original) The method of claim 46, wherein the subject has or is at risk of developing a cancer.
- 54. (Original) The method of claim 46, further comprising administering an antigen to the subject.
- 55. (Original). The method of claim 53, wherein the antigen is selected from the group consisting of a microbial antigen, a cancer antigen, a self antigen, and an allergen.
- 56. (Original) The method of claim 54, wherein the microbial antigen is selected from the group consisting of a bacterial antigen, a viral antigen, a fungal antigen, and a parasitic antigen.
- 57. (Original) The method of claim 55, wherein the antigen is derived from a microorganism selected from the group consisting of herpesviridae, retroviridae, orthomyroviridae, toxoplasma, haemophilus, campylobacter, clostridium, E.coli, and staphylococcus.
- 58. (Original) The method of claim 46, wherein the immune response is an antigenspecific immune response.

59.-63. (Cancelled)

64. (Original) The method of claim 46, further comprising administering to the subject a second therapeutic agent.

Serial No.: 10/613,228 - 7 -

Conf. No.: 4680

65. (Original) The method of claim 64, wherein the second therapeutic agent is an antimicrobial agent.

66. (Original) The method of claim 65, wherein the anti-microbial agent is selected from the group consisting of an anti-bacterial agent, an anti-viral agent, an anti-fungal agent, and an anti-parasite agent.

67.-70. (Cancelled)

- 71. (Original) The method of claim 46, wherein the immunostimulatory nucleic acid has a nucleotide backbone which includes at least one backbone modification.
- 72. (Original) The method of claim 71, wherein the backbone modification is a phosphorothioate modification.
 - 73. (Original) The method of claim 71, wherein the nucleotide backbone is chimeric.
- 74. (Original) The method of claim 71, wherein the nucleotide backbone is entirely modified.

75.-76. (Cancelled)

- 77. (Original) The method of claim 46, wherein the immunostimulatory nucleic acid is administered orally.
- 78. (Original) The method of claim 46, wherein the immunostimulatory nucleic acid is administered locally.
- 79. (Original) The method of cl aim 46, wherein the immunostimulatory nucleic acid is administered parenterally.

Serial No.: 10/613,228 - 8 -

Conf. No.: 4680

80. (Original) The method of claim 46, wherein the immunostimulatory nucleic acid is administered in a sustained release device.

81. (Original) The method of claim 46, wherein the immunostimulatory nucleic acid is administered to a mucosal surface.

82.-83. (Cancelled)

84. (Original) The method of claim 81, wherein the mucosal surface is selected from the group consisting of an oral, nasal, rectal, vaginal, and ocular surface.

85. (Original) The method of claim 46, further comprising isolating an immune cell from the subject, contacting the immune cell with an effective amount to activate the immune cell of the immunostimulatory nucleic acid and re-administering the activated immune cell to the subject.

86.-88. (Cancelled)

89. (Original) The method of claim 46, wherein the subject is a human.

90. (Original) The method of claim 46, wherein the subject is selected from the group consisting of a dog, cat, horse, cow, pig, sheep, goat, chicken, monkey and fish.

91.-94. (Cancelled)

95. (Previously Presented) The method of claim 53, wherein the cancer is selected from the group consisting of biliary tract cancer; bone cancer; brain and CNS cancer; breast cancer; cervical cancer; choriocarcinoma; colon cancer; connective tissue cancer; endometrial cancer; esophageal cancer; eye cancer; gastric cancer; Hodgkin's lymphoma; intraepithelial neoplasms; larynx cancer; lymphomas; liver cancer; lung cancer (e.g. small cell and non-small cell);

Serial No.: 10/613,228 - 9 -

Conf. No.: 4680

melanoma; neuroblastomas; oral cavity cancer; ovarian cancer; pancreas cancer; prostate cancer; rectal cancer; sarcomas; skin cancer; testicular cancer; thyroid cancer; and renal cancer.

96. (Original) The method of claim 46, further comprising administering an antibody specific for a cell surface antigen, and wherein the immune response results in antigen dependent cellular cytotoxicity (ADCC).

97. (Cancelled)

98. (Currently Amended) A method for inducing an innate immune response, comprising administering to the subject an the immunostimulatory nucleic acid of claim 1 in an amount effective for activating an innate immune response, wherein the immunostimulatory nucleic acid has a nucleotide sequence comprising SEQ ID NO:1.

99. (Cancelled)